



CRMG Company Profile

& RESEARCH EXPERIENCE

Present Administrative Staff

Medical Director/Principal Investigator, CRMG, Inc.	Dr. Javier Sosa Faria
President of Operations, CRMG, Inc.	Mario Franceschi
Vice- President, CRMG, Inc.	Yanira Franceschi
Sr. Coordinator, CRMG, Inc	Jailine Fraticelli

Contact Numbers

- Administration Service (787) 955-0800
- Accounting & Billing (787) 718-3130
- Operations & Clinical (787) 354-4420

Research Site-Affiliations & Medical Specialties

- Main Administrative Office: Urb. Jardines de San Domingo Calle 5 A-18 Juana Juana Diaz PR 00795

Medical facilities	Number of Professionals
Hospitals	2
Outpatient Hospital Clinics	6
Dialysis Centers	2
Infusion Centers	2
Emergency Room	8
Medical Offices	14
Oncology Infusion Centers	2

CRMG Medical Specialty	Number of Professionals
Nephrologist	3 New
Cardiologist	2
Internal Medicine	2
Pediatrician	2
Otolaryngologist	1
Pulmonologist	1
Gastroenterologist	2
General Practitioner	4
Oncologist	2

Company Work & Research Experience

- **(Motif Biosciences Inc.)** - A Phase 3, randomized, double-blinded, multicenter study to evaluate the safety and efficacy of intravenous Iclaprim versus Vancomycin in the treatment of acute bacterial skin and skin structure infections suspected or confirmed to be due to Gram-positive pathogens. (Revive 1)
- **(Gliknik Inc, Company)** - A Phase 2 Randomized, Double-Blind, Placebo-Controlled Clinical Trial to Determine the Safety and Efficacy of GL-0817 (with Cyclophosphamide) for the Prevention of Recurrence in HLA-A2+Patients with High-Risk Squamous Cell Carcinoma of the Oral Cavity.
- **(Motif Biosciences Inc.)** - A Phase 3, randomized, double-blind, multicenter study to evaluate the safety and efficacy of intravenous iclaprim versus vancomycin in the treatment of acute bacterial skin structure infections suspected or confirmed to be due to Gram-positive pathogens.
- **(Kyowa Pharmaceutical)** - A Phase 2, Multicenter, Randomized, Double-blind, Placebo-controlled Multiple Ascending Dose Study (Induction Therapy and Long-term Extension Therapy) of an Anti-OX40 Monoclonal Antibody (KHK4083) in Subjects with Moderately Active Ulcerative Colitis.
- **(Boehringer Ingelheim Pharmaceutical)** - Randomized, double-blind, placebo-controlled trial to investigate the safety, tolerability, pharmacokinetics and pharmacodynamics of 4 multiple rising oral doses of BI 685509 over 28 days in male and female patients with diabetic nephropathy
- **(Factor Therapeutic Ltd)** - A Prospective, Multi-Center, Double-Blind, Randomized, Placebo-Controlled Trial Comparing Two Doses of VF001-DP to Placebo as an Adjunct to Standard Care (SC) in Patients with Chronic Venous Leg Ulcers (VLUs)
- **(Emergent BioSolutions)** - A Randomized, Double-Blind, Placebo-Controlled Dose Ranging Study Evaluating the Safety, Pharmacokinetics and Clinical Benefit of FLU-IGIV in Hospitalized Patients with Serious Influenza A Infection
- **(Romark Global Pharma LCC.)** - A Phase III, Randomized, Double-Blind, Placebo Controlled Trial to Evaluate the Efficacy and Safety of Nitazoxanide in the Treatment of Uncomplicated Influenza.
- **(Romark Global Pharma LCC.)** - A Phase 3, Randomized, Double-Blind, Placebo Controlled Trial to Evaluate the Efficacy and Safety of Nitazoxanide in the Treatment of Colds due to Enterovirus/Rhinovirus Infection.
- **(Serena Group)** - A Randomized Controlled Multicenter Trial, Examining the Effect of Natrox Oxygen Wound Therapy on the Healing Rate of Chronic Diabetic Foot Ulcers (NOW.T-001)
- **(Serena Group)** Efficacy of TR 987, beta-1,3-1,6-D-glucan, in the treatment of chronic venous insufficiency ulcers: a two-arm, double blind, placebo controlled, randomized controlled trial. (BG001)

- **(Eli Lilly & Company)** - A Phase 3, Multicenter, Randomized, Double-Blind, Placebo- and ActiveControlled, Treat-Through Study to Evaluate the Efficacy and Safety of Mirikizumab in Patients With Moderately to Severely Active Crohn's Disease
- **(Serena Group)** - A Randomized Controlled Double-Blind Multi-Center Clinical Trial Evaluating Remote Ischemic Conditioning and Standard of Care vs Sham Therapy and Standard Care in the Healing of Diabetic Foot Ulcers
- **(Serena Group)** - A Randomized Controlled Multicenter Clinical Trial, Evaluating the Efficacy of Dual Layer Amniotic Membrane (Artacent®) and Standard of Care versus Standard of Care alone in the healing Chronic Diabetic Foot Ulcers
- **(Romark Global Pharma LCC.)** - A Randomized, Double-Blind, Placebo Controlled, Trial to Evaluate the Efficacy and Safety of Nitazoxanide (NTZ) For Post-Exposure Prophylaxis Of COVID-19 And Other Viral Respiratory Illnesses (VRI) In Healthcare Workers
- **(Romark Global Pharma LCC.)** - Phase 3, Randomized, Double-Blind, Placebo Controlled, Trial to Evaluate the Efficacy and Safety of Nitazoxanide In the Treatment Of Mild Or Moderate COVID-19
- **(CalciMedcia Biotechnology Company)** - A Randomized Double Blind, Placebo-Controlled Study of Auxora for the Treatment of Severe COVID-19 Pneumonia (CARDEA)
- **(Bristol-Myers Squibb)** - Phase 2, Randomized, Double-Blind Placebo Controlled Study of Intravenous Abatacept in the Treatment of Hospitalized COVID-19 Participants with Respiratory Compromise
- **(Pfizer Pharmaceutical)** - A Phase 2/3, randomized, double-blind, placebo-controlled study to evaluate the efficacy, safety and pharmacokinetics of orally administered PF 07321332/ritonavir in non-hospitalized adult participants with COVID-19 at low risk of progressing to severe disease
- **(Pfizer Pharmaceutical)** - A Phase 2/3, randomized, double-blind, placebo-controlled study to evaluate the efficacy, safety and pharmacokinetics of orally administered PF-07321332/ritonavir in non-hospitalized adult participants with COVID-19 at high risk of progressing to severe disease
- **(Pfizer Pharmaceutical)** - A Phase 2/3, randomized, double-blind, placebo-controlled study to evaluate the efficacy, safety, tolerability and pharmacokinetics of orally administered PF-07321332 in preventing SARS-CoV-2 infection in household contacts of individuals infected with SARS-CoV-2.
- **(Eli Lilly & Company)** - A Phase 2 Study of Once Daily LY3502970 Compared with Placebo and Once Weekly Dulaglutide in Participants with Type 2 Diabetes Mellitus
- **(AstraZeneca Pharmaceutical)** - A Randomized, Double-Blind, Double Dummy, Parallel Group, Multicenter 24 to 52 Week Variable Length Study to Assess the Efficacy and Safety of Budesonide,

Glycopyrronium, and Formoterol Fumarate Metered Dose Inhaler (MDI) Relative to Budesonide and Formoterol Fumarate MDI and Symbicort® Pressurized MDI in Adult and Adolescent Participants with Inadequately Controlled Asthma (LOGOS)